

IMMUNIZATION RESOURCES

FOR PARENTS &
PARENTS-TO-BE
2023



Dear Parent or Parent-to-Be,

This booklet contains helpful information on vaccines for you and your family.

Vaccines save lives. They protect against serious diseases like COVID-19, measles, flu, whooping cough, and chickenpox. Vaccinating children against certain diseases is one important way to help them stay healthy.

WIC is dedicated to partnering with other services to support your family's whole health, including by promoting immunizations. WIC staff can help you keep your child up-to-date on their vaccines -- all you need to do is either track your child's electronic vaccination records in your state's immunization information system (IIS) or keep a paper copy. If your child is behind on vaccinations, WIC staff will suggest a place you can go to get your child the vaccines they need.

Most health insurance plans cover the cost of vaccines, but you may want to check with your insurance provider before going to the doctor. If your child does not have health insurance, is Medicaid-eligible, or is under-insured, they may be able to receive free vaccines at their doctor's office or a health center through the federal Vaccines for Children program (VFC). For more information on VFC and how to pay for your family's vaccines, visit vaccinateyourfamily.org/vfc.

Adults, especially pregnant people, need vaccines too. By getting yourself vaccinated while pregnant you are helping to protect BOTH you and your baby from serious diseases such as flu, whooping cough, and COVID-19. For more information and for answers to questions you may have about vaccines for your family, please visit vaccinateyourfamily.org.

This booklet was created by Vaccinate Your Family and is supported through funding provided by The Centers for Disease Control and Prevention.

2023 Recommended Immunizations for Children from Birth Through 6 Years Old

VACCINE	Birth	1 MONTH	2 MONTHS	4 MONTHS	6 MONTHS	12 MONTHS	15 MONTHS	18 MONTHS	19-23 MONTHS	2-3 YEARS	4-6 YEARS
HepB Hepatitis B	HepB	HepB			HepB						
RV* Rotavirus			RV	RV	RV*						
DTaP Diphtheria, Pertussis, & Tetanus			DTaP	DTaP	DTaP		DTaP				DTaP
Hib* <i>Haemophilus influenzae</i> type b			Hib	Hib	Hib*	Hib					
PCV13, PCV15 Pneumococcal disease			PCV	PCV	PCV	PCV					
IPV Polio			IPV	IPV	IPV						IPV
COVID-19** Coronavirus disease 2019					COVID-19**						
Flu† Influenza					Flu (One or Two Doses Yearly)†						
MMR Measles, Mumps, & Rubella						MMR					MMR
Varicella Chickenpox						Varicella					Varicella
HepA[‡] Hepatitis A						HepA [‡]		HepA [‡]			

FOOTNOTES

RV* **Hib***
Administering a third dose at age 6 months depends on the brand of Hib or rotavirus vaccine used for previous dose.

COVID-19** Number of doses recommended depends on your child's age and type of COVID-19 vaccine used.

Flu† Two doses given at least 4 weeks apart are recommended for children age 6 months through 8 years of age who are getting an influenza (flu) vaccine for the first time and for some other children in this age group.

HepA[‡] Two doses of Hep A vaccine are needed for lasting protection. The 2 doses should be given between age 12 and 23 months. Both doses should be separated by at least 6 months. Children 2 years and older who have not received 2 doses of Hep A should complete the series.

ADDITIONAL INFORMATION

1. If your child misses a shot recommended for their age, talk to your child's doctor as soon as possible to see when the missed shot can be given.

2. If your child has any medical conditions that put them at risk for infection (e.g., sickle cell, HIV infection, cochlear implants) or is traveling outside the United States, talk to your child's doctor about additional vaccines that they may need.

Talk with your child's doctor if you have questions about any shot recommended for your child.



U.S. Department of Health and Human Services
Centers for Disease Control and Prevention

FOR MORE INFORMATION

Call toll-free: **1-800-CDC-INFO (1-800-232-4636)**
Or visit: **cdc.gov/vaccines/parents**



American Academy of Pediatrics



DEDICATED TO THE HEALTH OF ALL CHILDREN™

Diseases and the Vaccines that Prevent Them

BIRTH–6 YEARS OLD

DISEASE	VACCINE	DISEASE SPREAD BY	DISEASE SYMPTOMS	DISEASE COMPLICATIONS
Hepatitis B	HepB vaccine protects against hepatitis B.	Contact with blood or body fluids	May be no symptoms, fever, headache, weakness, vomiting, jaundice (yellowing of skin and eyes), joint pain	Chronic liver infection, liver failure, liver cancer, death
Rotavirus	RV vaccine protects against rotavirus.	Through the mouth	Diarrhea, fever, vomiting	Severe diarrhea, dehydration, death
Diphtheria	DTaP* vaccine protects against diphtheria.	Air, direct contact	Sore throat, mild fever, weakness, swollen glands in neck	Swelling of the heart muscle, heart failure, coma, paralysis, death
Pertussis (whooping cough)	DTaP* vaccine protects against pertussis (whooping cough).	Air, direct contact	Severe cough, runny nose, apnea (a pause in breathing in infants)	Pneumonia (infection in the lungs), death
Tetanus	DTaP* vaccine protects against tetanus.	Exposure through cuts in skin	Stiffness in neck and abdominal muscles, difficulty swallowing, muscle spasms, fever	Broken bones, breathing difficulty, death
Haemophilus influenzae type b (Hib)	Hib vaccine protects against <i>Haemophilus influenzae</i> type b.	Air, direct contact	May be no symptoms unless bacteria enter the blood	Meningitis (infection of the covering around the brain and spinal cord), intellectual disability, epiglottitis (life-threatening infection that can block the windpipe and lead to serious breathing problems), pneumonia (infection in the lungs), death
Pneumococcal disease (PCV13, PCV15)	PCV vaccine protects against pneumococcal disease.	Air, direct contact	May be no symptoms, pneumonia (infection in the lungs)	Bacteremia (blood infection), meningitis (infection of the covering around the brain and spinal cord), death
Polio	IPV vaccine protects against polio.	Air, direct contact, through the mouth	May be no symptoms, sore throat, fever, nausea, headache	Paralysis, death
Coronavirus disease 2019 (COVID-19)	COVID-19 vaccine protects against severe complications from coronavirus disease 2019.	Air, direct contact	May be no symptoms, fever, muscle aches, sore throat, cough, runny nose, diarrhea, vomiting, new loss of taste or smell	Pneumonia (infection in the lungs), respiratory failure, blood clots, bleeding disorder, injury to liver, heart or kidney, multi-system inflammatory syndrome, post-COVID syndrome, death
Influenza (Flu)	Flu vaccine protects against influenza.	Air, direct contact	Fever, muscle pain, sore throat, cough, extreme fatigue	Pneumonia (infection in the lungs), bronchitis, sinus infections, ear infections, death
Measles	MMR** vaccine protects against measles.	Air, direct contact	Rash, fever, cough, runny nose, pink eye	Encephalitis (brain swelling), pneumonia (infection in the lungs), death
Mumps	MMR** vaccine protects against mumps.	Air, direct contact	Swollen salivary glands (under the jaw), fever, headache, tiredness, muscle pain	Meningitis (infection of the covering around the brain and spinal cord), encephalitis (brain swelling), inflammation of testicles or ovaries, deafness, death
Rubella	MMR** vaccine protects against rubella.	Air, direct contact	Sometimes rash, fever, swollen lymph nodes	Very serious in pregnant women—can lead to miscarriage, stillbirth, premature delivery, birth defects
Chickenpox	Varicella vaccine protects against chickenpox.	Air, direct contact	Rash, tiredness, headache, fever	Infected blisters, bleeding disorders, encephalitis (brain swelling), pneumonia (infection in the lungs), death
Hepatitis A	HepA vaccine protects against hepatitis A.	Direct contact, contaminated food or water	May be no symptoms, fever, stomach pain, loss of appetite, fatigue, vomiting, jaundice (yellowing of skin and eyes), dark urine	Liver failure, arthralgia (joint pain), kidney, pancreatic and blood disorders, death

DTaP* DTaP combines protection against diphtheria, tetanus, and pertussis.

MMR** MMR combines protection against measles, mumps, and rubella.

Last updated December 2022 • CS322257-A


Immunizations and Developmental Milestones for Your Child from Birth Through 6 Years Old

Child's Name _____

Birth Date _____

		Birth	1 MONTH	2 MONTHS	4 MONTHS	6 MONTHS
Recommended Immunizations	Hepatitis B	<input type="radio"/> HepB	<input type="radio"/> HepB ¹			<input type="radio"/> HepB
	Rotavirus			<input type="radio"/> RV	<input type="radio"/> RV	<input type="radio"/> RV ²
	Diphtheria, Tetanus, Pertussis			<input type="radio"/> DTaP	<input type="radio"/> DTaP	<input type="radio"/> DTaP
	<i>Haemophilus influenzae</i> type b			<input type="radio"/> Hib	<input type="radio"/> Hib	<input type="radio"/> Hib
	Pneumococcal			<input type="radio"/> PCV	<input type="radio"/> PCV	<input type="radio"/> PCV
	Inactivated Poliovirus			<input type="radio"/> IPV	<input type="radio"/> IPV	<input type="radio"/> IPV
	Influenza (Flu)					<input type="radio"/> Influenza, first dose ³ <input type="radio"/> second dose
Milestones*	<p>Milestones should be achieved by the age indicated.</p> <p>Talk to your child's doctor about age-appropriate milestones if your child was born prematurely.</p> <ul style="list-style-type: none"> <input type="radio"/> Recognizes caregiver's voice <input type="radio"/> Turns head toward breast or bottle <input type="radio"/> Communicates through body language, fussing or crying, alert and engaged <input type="radio"/> Startles to loud sounds 	<ul style="list-style-type: none"> <input type="radio"/> Starts to smile <input type="radio"/> Raises head when on tummy <input type="radio"/> Calms down when rocked, cradled or sung to <input type="radio"/> Pays attention to faces 	<ul style="list-style-type: none"> <input type="radio"/> Begins to smile at people <input type="radio"/> Coos, makes gurgling sounds <input type="radio"/> Begins to follow things with eyes <input type="radio"/> Can hold head up 	<ul style="list-style-type: none"> <input type="radio"/> Babbles with expression <input type="radio"/> Likes to play with people <input type="radio"/> Reaches for toy with one hand <input type="radio"/> Brings hands to mouth <input type="radio"/> Responds to affection <input type="radio"/> Holds head steady, unsupported 	<ul style="list-style-type: none"> <input type="radio"/> Knows familiar faces <input type="radio"/> Responds to own name <input type="radio"/> Brings things to mouth <input type="radio"/> Rolls over in both directions <input type="radio"/> Strings vowels together when babbling ("ah", "eh", "oh") 	
Growth	At each well child visit, enter date, length, weight, and percentile information to keep track of your child's progress.	_____	_____	_____	_____	_____
		WEIGHT / PERCENTILE	WEIGHT / PERCENTILE	WEIGHT / PERCENTILE	WEIGHT / PERCENTILE	WEIGHT / PERCENTILE
		LENGTH / PERCENTILE	LENGTH / PERCENTILE	LENGTH / PERCENTILE	LENGTH / PERCENTILE	LENGTH / PERCENTILE
		_____	_____	_____	_____	_____
		HEAD CIRCUMFERENCE	HEAD CIRCUMFERENCE	HEAD CIRCUMFERENCE	HEAD CIRCUMFERENCE	HEAD CIRCUMFERENCE

continues on back page

 Shaded boxes indicate the vaccine can be given during shown age range.

VISIT DATE

VISIT DATE

VISIT DATE

VISIT DATE

VISIT DATE

¹ The second dose of HepB may be given either at the 1 month or 2 month visit.

² A third dose of rotavirus vaccine is only needed for RotaTeq.

³ Two doses given at least four weeks apart are recommended for children aged 6 months through 8 years of age who are getting a flu vaccine for the first time and for some other children in this age group.

* Milestones adapted from *Caring for your baby and young child: Birth to age 5*, Fifth Edition, edited by Steven Shelov and Tanya Remer Altmann © 1991, 1993, 1998, 2004, 2009 by the American Academy of Pediatrics and Bright Futures: Guidelines for health supervision of infants, children, and adolescents, Third Edition, edited by Joseph Hagan, Jr., Judith S. Shaw, and Paula M. Duncan, 2008, Elk Grove Village, IL: American Academy of Pediatrics.

This is not an exhaustive list of milestones from 0-6 years. See more at www.cdc.gov/milestones

If your child has any medical conditions that put him at risk for infections or is traveling outside the United States, talk to your child's doctor about additional vaccines that he may need.



U.S. Department of Health and Human Services
Centers for Disease Control and Prevention




American Academy of Pediatrics
DEDICATED TO THE HEALTH OF ALL CHILDREN™

Immunizations and Developmental Milestones for Your Child from Birth Through 6 Years Old

Child's Name _____

Birth Date _____

		12 MONTHS	15 MONTHS	18 MONTHS	19 23 MONTHS	2-3 YEARS	4-6 YEARS
Recommended Immunizations	Hepatitis B	<input type="radio"/> HepB (Final dose administered between 6 and 18 months)					
	Diphtheria, Tetanus, Pertussis		<input type="radio"/> DTaP				<input type="radio"/> DTaP
	Haemophilus influenzae type b	<input type="radio"/> Hib					
	Pneumococcal	<input type="radio"/> PCV					
	Inactivated Poliovirus	<input type="radio"/> IPV (Third dose administered between 6 and 18 months)					<input type="radio"/> IPV
	Influenza (Flu)	<input type="radio"/> Influenza , first dose ² <input type="radio"/> second dose (if needed)				Age 2 3 <input type="radio"/> <input type="radio"/> Influenza , first dose ² <input type="radio"/> <input type="radio"/> second dose (if needed)	Age 4 5 6 <input type="radio"/> <input type="radio"/> <input type="radio"/> Influenza , first dose ² <input type="radio"/> <input type="radio"/> <input type="radio"/> second dose (if needed)
	Measles, Mumps, Rubella	<input type="radio"/> MMR					<input type="radio"/> MMR
	Varicella	<input type="radio"/> Varicella					<input type="radio"/> Varicella
Hepatitis A	<input type="radio"/> <input type="radio"/> Hep A³						
Milestones*	Milestones should be achieved by the age indicated. Talk to your child's doctor about age-appropriate milestones if your child was born prematurely.	<input type="radio"/> Cries when mom or dad leaves <input type="radio"/> Says "mama" and "dada" <input type="radio"/> Copies gestures (for example, waves "bye bye") <input type="radio"/> May stand alone <input type="radio"/> Looks at right picture or thing when named	<input type="radio"/> Imitates what you are doing <input type="radio"/> Drinks from a cup <input type="radio"/> Scribbles on his own <input type="radio"/> Walks well <input type="radio"/> Says a couple of words other than "mama" and "dada"	<input type="radio"/> Points to show others something interesting <input type="radio"/> Says several single words <input type="radio"/> Points to one body part <input type="radio"/> May walk up steps and run	<input type="radio"/> Plays mainly beside other children <input type="radio"/> Follows two-step commands <input type="radio"/> Plays simple make-believe games <input type="radio"/> Throws ball overhand	<input type="radio"/> Can name most familiar things <input type="radio"/> Shows affection for friends without prompting <input type="radio"/> Turns book pages one at a time <input type="radio"/> Kicks a ball	<input type="radio"/> Speaks very clearly <input type="radio"/> Tells stories <input type="radio"/> Can print some letters or numbers <input type="radio"/> Hops; may be able to skip <input type="radio"/> Enjoys playing with other children
	Growth	At each well child visit, enter date, length, weight, and percentile information to keep track of your child's progress.	_____ WEIGHT / PERCENTILE _____ LENGTH / PERCENTILE _____ HEAD CIRCUMFERENCE	_____ WEIGHT / PERCENTILE _____ LENGTH / PERCENTILE _____ HEAD CIRCUMFERENCE	_____ WEIGHT / PERCENTILE _____ LENGTH / PERCENTILE _____ HEAD CIRCUMFERENCE	_____ WEIGHT / PERCENTILE _____ LENGTH / PERCENTILE _____ HEAD CIRCUMFERENCE	_____ WEIGHT _____ HEIGHT _____ BMI

 Shaded boxes indicate the vaccine can be given during shown age range.

VISIT DATE

VISIT DATE

VISIT DATE

VISIT DATE

VISIT DATE

VISIT DATE

² Two doses given at least four weeks apart are recommended for children aged 6 months through 8 years of age who are getting a flu vaccine for the first time and for some other children in this age group.

³ Two doses of HepA vaccine are needed for lasting protection. The first dose of HepA vaccine should be given between 12 months and 23 months of age. The second dose should be given 6 months after the first dose. HepA vaccination may be given to any child 12 months and older to protect against HepA. Children and adolescents who did not receive the HepA vaccine and are at high-risk, should be vaccinated against HepA.

* Milestones adapted from AAP *Bright Futures: Guidelines for Health Supervision of Infants, Children, and Adolescents* Third Edition.

This is not an exhaustive list of milestones from 0-6 years. See more at www.cdc.gov/Milestones and download the Milestone Tracker App at www.cdc.gov/MilestoneTracker

If your child has any medical conditions that put him at risk for infections or is traveling outside the United States, talk to your child's doctor about additional vaccines that he may need.



U.S. Department of Health and Human Services
Centers for Disease Control and Prevention




American Academy of Pediatrics



DEDICATED TO THE HEALTH OF ALL CHILDREN™

2023 Recommended Immunizations for Children 7–18 Years Old



KEY



Indicates when the vaccine is recommended for all children unless your doctor tells you that your child cannot safely receive the vaccine.



Indicates the vaccine series can begin at this age.



Indicates the vaccine **should** be given if a child is catching up on missed vaccines. A vaccine series does not need to be restarted, regardless of the time that has elapsed between doses.



Indicates children not at increased risk **may** get the vaccine if they wish after speaking to a provider.

ADDITIONAL INFORMATION

1. If your child misses a shot recommended for their age, talk to your child's doctor as soon as possible to see when the missed shot can be given.
2. If your child has any medical conditions that put them at risk for infection or is traveling outside the United States, talk to your child's doctor about additional vaccines that they may need.

Talk with your child's doctor if you have questions about any shot recommended for your child.

FOOTNOTES

COVID-19* Number of doses recommended depends on your child's age and type of COVID-19 vaccine used.

Flu** Two doses given at least 4 weeks apart are recommended for children age 6 months through 8 years of age who are getting an influenza (flu) vaccine for the first time and for some other children in this age group.

HPV+ Ages 11 through 12 years old should get a 2-shot series separated by 6 to 12 months. The series can begin at 9 years old. A 3-shot series is recommended for those with weakened immune systems and those who start the series after their 15th birthday.

*Originally recommended age ranges for missed childhood vaccinations: 2-dose series of **MMR** at 12–15 months and 4–6 years; 2-dose series of **Varicella** at 12–15 months and 4–6 years; 2-dose series of **HepA** (minimum interval: 6 months) at age 12–23 months; 3-dose series of **HepB** at birth, 1–2 months, and 6–18 months; and 4-dose series of **Polio** at 2 months, 4 months, 6–18 months, and 4–6 years.



U.S. Department of Health and Human Services
Centers for Disease Control and Prevention

FOR MORE INFORMATION

Call toll-free: **1-800-CDC-INFO (1-800-232-4636)**
Or visit: cdc.gov/vaccines/parents




American Academy of Pediatrics



DEDICATED TO THE HEALTH OF ALL CHILDREN™

COVID-19 IMPACTS KIDS, TOO!



Nearly **200,000 children** have been hospitalized with COVID-19

While most children with COVID-19 have mild symptoms or no symptoms at all, children can – and some do – get severely ill from COVID. Those that get very sick from COVID could need to be hospitalized. In rare cases, they might die.

COVID-19 VACCINES ARE THE BEST PROTECTION AGAINST THE VIRUS THAT CAUSES COVID.

Vaccines are now recommended for babies and children 6 months and older. The vaccine provides protection against severe disease, hospitalization, and death.

To see if your children are up to date on their vaccines, look at the CDC's immunization schedule and talk to your healthcare provider.

Infant Immunizations FAQs

It's natural you have questions about your child's vaccines. Read answers to common questions to learn more about vaccine safety, the recommended schedule, how vaccines protect your child from 14 diseases by age two, and more. CDC regularly updates this document to ensure frequently asked questions from parents are answered with the most current information.

Q: Are vaccines safe?

A: Yes. Vaccines are very safe. The United States' long-standing vaccine safety system ensures that vaccines are as safe as possible. Currently, the United States has the safest vaccine supply in its history. Millions of children safely receive vaccines each year. The most common side effects are typically very mild, such as pain or swelling at the injection site.

Q: What are the side effects of the vaccines? How do I treat them?

A: Vaccines, like any medication, may cause some side effects. **Most of these side effects are very minor, like soreness where the shot was given, fussiness, or a low-grade fever.** These side effects typically only last a couple of days and are treatable. For example, you can apply a cool, wet washcloth on the sore area to ease discomfort.

Serious reactions are very rare. However, if your child experiences any reactions that concern you, call the doctor's office.

Q: What are the risks and benefits of vaccines?

A: Vaccines can prevent infectious diseases that once killed or harmed many infants, children, and adults. Without vaccines, your child is at risk for getting seriously ill and suffering pain, disability, and even death from diseases like measles and whooping cough. The main risks associated with getting vaccines are side effects, which are almost always mild (redness and swelling at the injection site) and go away within a few days. Serious side effects after vaccination, such as a severe allergic reaction, are very rare and doctors and clinic staff are trained to deal with them. **The disease-prevention benefits of getting vaccines are much greater than the possible side effects for almost all children.** The only exceptions to this are cases in which a child has a serious chronic medical condition like cancer or a disease that weakens the immune system, or has had a severe allergic reaction to a previous vaccine dose.

Q: Is there a link between vaccines and autism?

A: No. Scientific studies and reviews continue to show no relationship between vaccines and autism.

Some people have suggested that thimerosal (a compound that contains mercury) in vaccines given to infants and young children might be a cause of autism. Others have suggested that the MMR (measles-mumps-rubella) vaccine may be linked to autism. However, numerous scientists and researchers have studied and continue to study the MMR vaccine and thimerosal, and they reach the same conclusion: there is no link between MMR vaccine or thimerosal and autism.

Q: Can vaccines overload my baby's immune system?

A: Vaccines do not overload the immune system. Every day, a healthy baby's immune system successfully fights off thousands of germs. Antigens are parts of germs that cause the body's immune system to go to work to build antibodies, which fight off diseases.

The antigens in vaccines come from the germs themselves, but the germs are weakened or killed so they cannot cause serious illness. **Even if babies receive several vaccinations in one day, vaccines contain only a tiny fraction of the antigens they encounter every day in their environment.** Vaccines give your child the antibodies they need to fight off serious vaccine-preventable diseases.



CDC recommends all children receive vaccines according to the recommended immunization schedule to protect them from 14 diseases by age two. Read below to get answers to 19 common questions about how vaccines benefit your child, the vaccine schedule, and more.

Q: Why are so many doses needed for each vaccine?

A: Getting every recommended dose of each vaccine provides your child with the best protection possible. Depending on the vaccine, your child will need more than one dose to build high enough immunity to prevent disease or to boost immunity that fades over time. Your child may also receive more than one dose to make sure they are protected if they did not get immunity from a first dose, or to protect them against germs that change over time, like flu. Every dose is important because each protects against infectious diseases that can be especially serious for infants and very young children.

Q: Why do vaccines start so early?

A: The recommended schedule protects infants and children by providing immunity early in life, before they come into contact with life-threatening diseases. Children receive immunization early because they are susceptible to diseases at a young age. The consequences of these diseases can be very serious, even life-threatening, for infants and young children.

Q: What do you think of delaying some vaccines or following a non-standard schedule?

A: Children do not receive any known benefits from following schedules that delay vaccines. Infants and young children who follow immunization schedules that spread out or leave out shots are at risk of developing diseases during the time you delay their shots. Some vaccine-preventable diseases remain common in the United States and children may be exposed to these diseases during the time they are not protected by vaccines, placing them at risk for a serious case of the disease that might cause hospitalization or death.

Infant Immunizations FAQs

Q: Haven't we gotten rid of most of these diseases in this country?

A: Some vaccine-preventable diseases, like pertussis (whooping cough) and chickenpox, remain common in the United States. On the other hand, other diseases vaccines prevent are no longer common in this country because of vaccines. **However, if we stopped vaccinating, the few cases we have in the United States could very quickly become tens or hundreds of thousands of cases.** Even though many serious vaccine-preventable diseases are uncommon in the United States, some are common in other parts of the world. Even if your family does not travel internationally, you could come into contact with international travelers anywhere in your community. Children who don't receive all vaccinations and are exposed to a disease can become seriously sick and spread it through a community.

Q: What are combination vaccines? Why are they used?

A: Combination vaccines protect your child against more than one disease with a single shot. They reduce the number of shots and office visits your child would need, which not only saves you time and money, but also is easier on your child.

Some common combination vaccines are Pediarix®, which combines DTap, Hep B, and IPV (polio), and ProQuad®, which combines MMR and varicella (chickenpox).

Q: Can't I just wait until my child goes to school to catch up on immunizations?

A: Before entering school, young children can be exposed to vaccine-preventable diseases from parents and other adults, brothers and sisters, on a plane, at child care, or even at the grocery store. Children under age 5 are especially susceptible to diseases because their immune systems have not built up the necessary defenses to fight infection. **Don't wait to protect your baby and risk getting these diseases when he or she needs protection now.**

Q: Why does my child need a chickenpox shot? Isn't it a mild disease?

A: Your child needs a chickenpox vaccine because chickenpox can actually be a serious disease. In many cases, children experience a mild case of chickenpox, but other children may have blisters that become infected. Others may develop pneumonia. There is no way to tell in advance how severe your child's symptoms will be.

Before vaccine was available, about 50 children died every year from chickenpox, and about 1 in 500 children who got chickenpox was hospitalized.

Q: My child is sick right now. Is it okay for her to still get shots?

A: Talk with your child's doctor, but children can usually get vaccinated even if they have a mild illness like a cold, earache, mild fever, or diarrhea. If the doctor says it is okay, your child can still get vaccinated.

Q: What are the ingredients in vaccines and what do they do?

A: Vaccines contain ingredients that cause the body to develop immunity. Vaccines also contain very small amounts of other ingredients. **All ingredients play necessary roles either in making the vaccine, or in ensuring that the final product is safe and effective.**

Q: Don't infants have natural immunity? Isn't natural immunity better than the kind from vaccines?

A: Babies may get some temporary immunity (protection) from mom during the last few weeks of pregnancy, but only for diseases to which

mom is immune. Breastfeeding may also protect your baby temporarily from minor infections, like colds. **These antibodies do not last long, leaving your baby vulnerable to disease.**

Natural immunity occurs when your child is exposed to a disease and becomes infected. It is true that natural immunity usually results in better immunity than vaccination, but the risks are much greater. A natural chickenpox infection may result in pneumonia, whereas the vaccine might only cause a sore arm for a couple of days.

Q: Can't I just wait to vaccinate my baby, since he isn't in child care, where he could be exposed to diseases?

A: No, even young children who are cared for at home can be exposed to vaccine preventable diseases, so it's important for them to get all their vaccines at the recommended ages. Children can catch these illnesses from any number of people or places, including from parents, brothers or sisters, visitors to their home, on playgrounds or even at the grocery store. Regardless of whether or not your baby is cared for outside the home, she comes in contact with people throughout the day, some of whom may be sick but not know it yet.

If someone has a vaccine preventable disease, they may not have symptoms or the symptoms may be mild, and they can end up spreading disease to babies or young children. Remember, many of these diseases can be especially dangerous to young children so it is safest to vaccinate your child at the recommended ages to protect her, whether or not she is in child care.

Q: Do I have to vaccinate my baby on schedule if I'm breastfeeding him?

A: Yes, even breastfed babies need to be protected with vaccines at the recommended ages. The immune system is not fully developed at birth, which puts newborns at greater risk for infections.

Breast milk provides important protection from some infections as your baby's immune system is developing. For example, babies who are breastfed have a lower risk of ear infections, respiratory tract infections, and diarrhea. However, breast milk does not protect children against all diseases. Even in breastfed infants, vaccines are the most effective way to prevent many diseases. Your baby needs the long-term protection that can only come from making sure he receives all his vaccines according to the CDC's recommended schedule.

Q: What's wrong with delaying some of my baby's vaccines if I'm planning to get them all eventually?

A: Young children have the highest risk of having a serious case of disease that could cause hospitalization or death. Delaying or spreading out vaccine doses leaves your child unprotected during the time when they need vaccine protection the most. For example, diseases such as Hib or pneumococcus almost always occur in the first 2 years of a baby's life. And some diseases, like Hepatitis B and whooping cough (pertussis), are more serious when babies get them at a younger age. Vaccinating your child according to the CDC's recommended immunization schedule means you can help protect him at a young age.

Q: I got the whooping cough and flu vaccines during my pregnancy. Why does my baby need these vaccines too?

A: The protection (antibodies) you passed to your baby before birth will give him some early protection against whooping cough and flu. However, these antibodies will only give him short-term protection. It is very important for your baby to get vaccines on time so he can start building his own protection against these serious diseases.



VACCINES DURING PREGNANCY

Getting vaccinated while pregnant protects both you AND your baby

Flu can pose serious health risks for pregnant people and newborns.

Getting the flu shot during pregnancy is a simple way to help protect yourself and your baby from the flu.



Flu shots are safe for you and your baby during any trimester of your pregnancy.

Protect yourself and your baby by getting whooping cough vaccines during the 3rd trimester of every pregnancy!



Whooping cough can be lethal for infants.

Vaccinating during pregnancy gives your baby antibodies that protect them until they can get their vaccine at 2 months.

Pregnant and recently-pregnant people are at higher risk of serious COVID-19 illness than those who are not pregnant.

Getting sick with COVID-19 during pregnancy may also put your baby at increased risk of stillbirth and preterm birth



For the best protection from COVID-19, pregnant, recently-pregnant & breastfeeding people should get the COVID-19 vaccine ASAP!

Surround Your Baby with Protection

Ask friends, family and caregivers who plan to visit your baby to be up-to-date on all of their vaccinations at least two weeks before meeting your newborn.

Learn more about vaccines during pregnancy at [vaccinateyourfamily.org](https://www.vaccinateyourfamily.org).

Keeping Track of Your Child's Vaccines: Q & A for Parents

What are Immunization Information Systems?

Immunization Information Systems (IIS), also known as immunization registries, are electronic systems that have information on the vaccines that were given to your child. Each state has its own IIS. **There is NO cost for having records in IIS.**

What are the benefits of having my child's vaccine record saved in an IIS?

- Helps your child's doctors and/or other vaccine providers keep track of the shots due for your child.
- Helps your doctor send you vaccine reminders for your child.
- Helps to make sure that your child doesn't miss any shots **OR** get too many shots.
- Allows you to quickly get a copy of your child's shot record from the doctor.
- **Makes sure your child has all of the vaccines needed to start daycare, school and/or camp, and for international travel.**

What information is in my state's IIS?

The vaccine records in each state's IIS is kept confidential, and only authorized users are able to access the information. The exact information stored in the IIS depends on which state (or city) you live in, but most contain:



- Patient's name (first, middle and last)
- Patient's birth date
- Patient's gender
- Patient's birth state/country
- Mother's name
- Types of shots given
- Dates the shots were given

Who do I contact to see if my child's shot record is in my state's IIS or if I want a copy of my child's shot record?

- Contact your doctor's office or other vaccine provider to ask if your child's shot record is in the IIS and request a copy
- Contact the IIS in your state or in the state where your child got their last shots to see if they have records in the IIS. **To find contact information for your state's IIS, visit www.cdc.gov/vaccines/programs/iis/contacts-locate-records.html#state.** A few states allow the public to directly access the IIS in order to print out their child's vaccine records.

HOW TO PAY FOR VACCINES

No one should skip vaccines because of their **cost**.



Use Vaccinate Your Family's online tool to help you find out how to pay for your family's vaccines.

1

Enter the person's health insurance status: private insurance, public insurance or no insurance

2

Enter the person's age



Check out the Paying for Vaccines Tool and helpful vaccine information on our website at

vaccinateyourfamily.org

Tips for a Less Stressful Shot Visit



Making the choice to vaccinate your child is vital for their health and well-being. Even so, getting shots can still be stressful for you and your little one. Fortunately, there are simple ways you can support your child before, during, and after shots.

Before Getting Shots

Come prepared! Take these steps before your child gets a shot to help make the immunization visit less stressful on you both.

- Read any vaccine materials you received from your child's health care professional and write down any questions you may have.
- Find your child's personal immunization record and bring it to your appointment. An up-to-date record tells your doctor exactly what shots your child has already received.
- Pack a favorite toy or book, and a blanket that your child uses regularly to comfort your child.

For older children

- Be honest with your child. Explain that shots can pinch or sting, but that it won't hurt for long.
- Engage other family members, especially older siblings, to support your child.
- Avoid telling scary stories or making threats about shots.

At the Doctor's Office

If you have questions about immunizations, ask your child's doctor or nurse. Before you leave the appointment, ask your child's doctor for advice on using non-aspirin pain reliever and other steps you can take at home to comfort your child.

Try these ideas for making the shots easier on your child.

- Distract and comfort your child by cuddling, singing, or talking softly.
- Smile and make eye contact with your child. Let your child know that everything is ok.
- Comfort your child with a favorite toy or book. A blanket that smells familiar will help your child feel more comfortable.
- Hold your child firmly on your lap, whenever possible.

Help children see vaccines as a good thing. Never threaten your child with shots, by saying, "If you misbehave I will have the nurse give you a shot." Instead, remind children that vaccines can keep them healthy.

Ways to soothe your baby:

- Swaddling
- Skin-to-skin contact
- Offering a sweet beverage, like juice (when the child is older than 6 months)
- Breastfeeding

Your health care professional may cool or numb the injection site to reduce the pain associated with your child's shots.

The Centers for Disease Control and Prevention (CDC), the American Academy of Family Physicians (AAFP), and the American Academy of Pediatrics (AAP) adapted this information from *Be There for Your Child during Shots*, California Department of Public Health Immunization Branch.

For older children

- Take deep breaths with your child to help “blow out” the pain.
- Point out interesting things in the room to help create distractions.
- Tell or read stories.
- Support your child if he or she cries. Never scold a child for not “being brave.”

Once your child has received all of the shots, be especially supportive. Hold, cuddle, and, for infants, breastfeed or offer a bottle. A soothing voice, combined with praise and hugs will help reassure your child that everything is ok.


After the Shots

Sometimes children experience mild reactions from vaccines, such as pain at the injection site, a rash or a fever. These reactions are normal and will soon go away. The following tips will help you identify and minimize mild side effects.

- Review any information your doctor gives you about the shots, especially the Vaccine Information Statements or other sheets that outline which side effects might be expected.
- Use a cool, wet cloth to reduce redness, soreness, and swelling in the place where the shot was given.
- Reduce any fever with a cool sponge bath. If your doctor approves, give non-aspirin pain reliever.
- Give your child lots of liquid. It’s normal for some children to eat less during the 24 hours after getting vaccines.
- Pay extra attention to your child for a few days. If you see something that concerns you, call your doctor.

*Remember to schedule your next visit!
Staying current with your child’s immunizations
provides the best protection against disease.*

*Take a moment to read the Vaccine Information
Sheet your health care professional gives you
during your visit. This sheet has helpful
information and describes possible side effects
your child may experience.*



Credible Organizations & Websites for Vaccine Information

Vaccinate Your Family (VYF)

vaccinateyourfamily.org
vaccinateyourfamily.org/covidfaq
vaccinateyourfamily.org/current-flu-
season

American Academy of Family Physicians (AAFP)

aafp.org & familydoctor.org

American Academy of Pediatrics (AAP)

aap.org & healthychildren.org

American College of Nurse-Midwives

midwife.org

American College of Obstetricians & Gynecologists (ACOG)

acog.org

Centers for Disease Control and Prevention (CDC)

cdc.gov/vaccines
cdc.gov/coronavirus

Cervivor

cervivor.org

Emily Stillman Foundation

foreveremily.org

Good Health WINS

goodhealthwins.org

Kimberly Coffey Foundation

kimberlycoffeyfoundation.org

Families Fighting Flu (FFF)

familiesfightingflu.org

HPV Vaccination Roundtable

hpvroundtable.org

Immunize.org

immunize.org & vaccineinformation.org

Institute for Vaccine Safety at Johns Hopkins – Bloomberg School of Public Health

vaccinesafety.edu

March of Dimes

marchofdimes.org

National Foundation for Infectious Diseases (NFID)

nfid.org

National Meningitis Association

nmaus.org

Meningitis B Action Project

meningitisbactionproject.org

Society for Maternal-Fetal Medicine

smfm.org/covidfamily

Vaccine Education Center at The Children's Hospital of Philadelphia (CHOP)

vaccine.chop.edu
COVIDVaccineAnswers.org

Voices for Vaccines (VFV)

nmaus.org



Check Out Our Website!


Find vaccine resources for the whole family!



Watch our YouTube series, "Vaccines Explained," to learn about vaccine safety and how vaccines move throughout the body!



Check out our Child and Teen Vaccine-Preventable Disease eBook for more information on each of the diseases vaccines can prevent.



Check out our Adult Vaccine-Preventable Disease eBooks for more information on the diseases vaccines can prevent.

Visit Today!

vaccinateyourfamily.org